

IN THE CLAIMS

1. (currently amended) A method of screening for therapeutic agents useful in the treatment of a disease selected from ~~hematological diseases, disorders of the peripheral and central nervous system, respiratory diseases,~~ cardiovascular diseases, COPD, asthma, and genito-urological disorders in a mammal, comprising the steps of

- i) contacting a test compound with an N-formyl peptide receptor like-1 (FPRL1) polypeptide comprising the amino acid sequence SEQ ID NO:2,
- ii) detecting binding of said test compound to said FPRL1 polypeptide, and
- iii) identifying a test compound which binds to said FPRL1 polypeptide as a potential therapeutic agent useful in the treatment of the disease.

2. (currently amended) A method of screening for therapeutic agents useful in the treatment of a disease selected from ~~hematological diseases, disorders of the peripheral and central nervous system, respiratory diseases,~~ cardiovascular diseases, COPD, asthma, and genito-urological disorders in a mammal, comprising the steps of

- i) determining the activity of a FPRL1 polypeptide comprising the amino acid sequence SEQ ID NO:2 in the presence and absence of said test compound, and
- ii) identifying the test compound as a potential therapeutic agent useful in the treatment of the disease if the activity of the FPRL1 polypeptide is inhibited in the presence but not the absence of the test compound.

3. (currently amended) A method of screening for therapeutic agents useful in the treatment of a disease selected from ~~hematological diseases, disorders of the peripheral and central nervous system, respiratory diseases,~~ cardiovascular diseases, COPD, asthma, and genito-urological disorders in a mammal, comprising the steps of

- i) determining the activity of a FPRL1 polypeptide comprising the amino acid sequence SEQ ID NO:2 at a certain concentration of a test compound,
- ii) determining the activity of a FPRL1 polypeptide in the presence of a compound known to be a regulator of a FPRL1 polypeptide, and
- iii) identifying the test compound as a potential therapeutic agent useful in the treatment of the disease if the activity of the FPRL1 polypeptide is inhibited in the presence of the test compound and the compound known to be a regulator.
4. (original) The method of claim 1, wherein the step of contacting is in or at the surface of a cell.
5. (previously presented) The method of claim 4, wherein the cell is *in vitro*.
6. (previously presented) The method of claim 1, wherein the contacting is in a cell-free system.
7. (previously presented) The method of claim 1, wherein the polypeptide is coupled to a detectable label.
8. (previously presented) The method of claim 1, wherein the test compound is coupled to a detectable label.
9. (previously presented) The method of claim 1, wherein the test compound displaces a ligand which is bound to the polypeptide before the step of contacting.
10. (previously presented) The method of claim 1, wherein the polypeptide is attached to a solid support.
11. (previously presented) The method of claim 1, wherein the compound is attached to a solid support.

12. (withdrawn) A method of screening for therapeutic agents useful in the treatment of a disease selected from ~~hematological diseases, cardiovascular diseases, disorders of the peripheral and central nervous system, respiratory diseases,~~ COPD, asthma, and genito-urological disorders in a mammal, comprising the steps of

- i) contacting a test compound with a FRPL1 polynucleotide encoding an FRPL1 polypeptide comprising the amino acid sequence SEQ ID NO:2,
- ii) detect binding of said test compound to said FRPL1 polynucleotide, and
- iii) identifying a test compound which binds to the FRPL1 polynucleotide as a potential therapeutic agent useful in the treatment of the disease.

13. (withdrawn) The method of claim 12 wherein the nucleic acid molecule is RNA.

14. (withdrawn) The method of claim 12 wherein the contacting step is in or at the surface of a cell.

15. (withdrawn) The method of claim 12 wherein the contacting step is in a cell-free system.

16. (withdrawn) The method of claim 12 wherein the polynucleotide is coupled to a detectable label.

17. (withdrawn) The method of claim 12 wherein the test compound is coupled to a detectable label.

18. (withdrawn) A method of diagnosing a disease selected from ~~hematological diseases, cardiovascular diseases, disorders of the peripheral and central nervous system, respiratory diseases,~~ COPD, asthma, and genito-urological disorders in a mammal, comprising the steps of

- i) determining the amount of a FRPL1 polynucleotide encoding an FRPL1 polypeptide comprising the amino acid sequence SEQ ID NO:2 in a sample taken from said mammal,

ii) determining the amount of FPRL1 polynucleotide in healthy and/or diseased mammals.

19-26. (canceled)